

DETERMINATION OF THE DIRECT CAPTURE CONTRIBUTION FOR $^{13}\text{N}(p,\gamma)^{14}\text{O}$ FROM THE $^{14}\text{O} \rightarrow ^{13}\text{N} + p$ ASYMPTOTIC NORMALIZATION COEFFICIENT

Xiaodong Tang^{a),b)}, A. Azhari^{b)}, Changbo Fu^{b)}, C.A. Gagliardi^{b)}, A.M. Mukhamedzhanov^{b)}, F. Pirlepesov^{b)}, L. Trache^{b)}, R.E. Tribble^{b)}, V. Burjan^{c)}, V. Kroha^{c)}, and B. F. Irgaziev^{d)}

^{a)} *Argonne National Laboratory, Argonne, IL 60439, USA*

^{b)} *Cyclotron Institute, Texas A&M University, College Station, TX 77843*

^{c)} *Institute of Nuclear Physics, Czech Academy of Sciences, Prague-Řež, Czech Republic*

^{d)} *Department of Physics, National University, Tashkent, Uzbekistan*

$^{13}\text{N}(p,\gamma)^{14}\text{O}$ is one of the key reactions which trigger the onset of the hot CNO cycle [1]. This transition occurs when the proton capture rate on ^{13}N is faster, due to increasing stellar temperature ($\geq 10^8$ K), than the ^{13}N β -decay rate [2]. The rate of this reaction is dominated by the resonant capture through the first excited state of ^{14}O ($E_r = 0.528$ MeV). However, through constructive interference direct capture below the resonance makes a non-negligible contribution to the reaction rate. We have determined this direct contribution by measuring the asymptotic normalization coefficient for $^{14}\text{O} \rightarrow ^{13}\text{N} + p$. In our experiment, an 11.8 MeV/nucleon ^{13}N radioactive beam was used to study the $^{14}\text{N}(^{13}\text{N}, ^{14}\text{O})^{13}\text{C}$ peripheral transfer reaction and the asymptotic normalization coefficient, $(C_{p_{1/2}}^{^{14}\text{O}})^2 = 29.0 \pm 4.3$ fm⁻¹, was extracted from the measured cross section. The radiative capture cross section was estimated using an R -matrix approach with the measured asymptotic normalization coefficient and the latest resonance parameters. We find the S factor for $^{13}\text{N}(p,\gamma)^{14}\text{O}$ to be larger than previous estimates [3]. Consequently the transition from the cold to hot CNO cycle for novae would be controlled by the slowest proton capture reaction $^{14}\text{N}(p,\gamma)^{15}\text{O}$.

[1] M. Wiescher, J. Gorres and H. Schatz, J. Phys. G **25**, R133 (1999).

[2] M.S. Smith *et al.*, Phys. Rev. C **47**, 2740 (1993).

[3] P. Decrock *et al.*, Phys. Rev. C **48**, 2057 (1993).